

On Singularity Formation Under Mean Curvature Flow

I.M.Sigal

Toronto

Joint work with Wenbin Kong, Zhou Gang,
and Dan Knopf

Also related work with
Dimitra Antonopoulou and Georgia Karali

Heraklion, May 2013

Mean Curvature Flow

The mean curvature flow is a family of hypersurfaces $M_t \subset \mathbb{R}^{d+1}$ whose smooth immersions $\psi(\cdot, t) : N \rightarrow M_t \subset \mathbb{R}^{d+1}$ satisfy the partial differential equation

$$(\partial_t \psi)^N = -H(\psi)$$

where $(\partial_t \psi)^N$ is the normal component of $\partial_t \psi$ and $H(x)$ is the mean curvature of M_t at a point $x \in M_t$.

Applications and Connections

- ▶ Material Science (interface motion between different materials or different phases).
- ▶ Image recognition.
- ▶ Connection to the Ricci flow.
- ▶ Topological classification of surfaces and submanifolds.

Some Key Works: Existence

- ▶ First mathematical treatment (using geometric measure theory): Brakke [1978];
- ▶ Short time existence: Brakke, Huisken, Evans and Spruck, Ilmanen, Ecker and Huisken [1991];
- ▶ Weak solutions: Evans and Spruck, Chen, Giga and Goto [1991];

Some Key Works: Singularities

The most interesting problem here is formation of singularities.

- ▶ Collapse of convex hypersurfaces: Huisken [1984], extensions: White [2000, 2003], Huisken and Sinestrari [2007-2009];
- ▶ Neckpinching for rotationally symmetric hypersurfaces: Grayson, Ecker, Huisken, M. Simon, Dziuk and Kawohl, Smoczyk, Altschuler, Angenent and Giga, Soner and Souganidis [1990-1995];
- ▶ MCF with surgery and topological classification of surfaces and submanifolds: Huisken and Sinestrari [2007-2009];
- ▶ Nature of the singular set: White [2000, 2003], Colding and Minicozzi [2012].

There are three explicit solutions of MCF:

- ▶ Collapsing Euclidean spheres with radii decreasing as $\sqrt{2d(t_* - t)}$;
- ▶ Collapsing Euclidean cylinders with radii decreasing as $\sqrt{2(d-1)(t_* - t)}$;

Conjecture [Huisken]: Generic sing. are spheres and cylinders.

Partial results: Huisken, White, Colding and Minicozzi

Results:

- ▶ Spherical collapse is asymptotically stable.
- ▶ Cylindrical collapse is unstable.
- ▶ Cylindrical collapse \Rightarrow neckpinching (stable and universal).

Stability of Shrinking Spheres

Theorem. (W. Kong-I.M.S.) Let a surface M_0 be close to S^d in H^s , $s > \frac{d}{2} + 1$. Then $\exists t_* < \infty$, s.t. MCF has a solution M_t for $0 \leq t < t_*$ and

- ▶ $M_t \rightarrow z_*$, for some z_* , as $t \rightarrow t_*$;
- ▶ M_t are defined by immersions of S^d , satisfying

$$\psi(\omega, t) = z(t) + \rho(\omega, t)\omega,$$

$$\rho(t) = \sqrt{\tau} \left(1 + O_{H^s}(\tau^\beta) \right),$$

with $\tau := 2d(t_* - t)$, $\alpha := \frac{1}{2}(d + \frac{1}{2} - \frac{1}{2d})$ and $\beta := \frac{1}{2}(1 - \frac{1}{2d})$.

Graphs over Cylinders

Our next result deals with initial conditions M_0 , which are graphs over $(d + 1)$ -dimensional cylinders C^{d+1} along the x_{d+2} -axis in \mathbb{R}^{d+2} ,

$$\psi_0(\omega, x) = (u_0(\omega, x)\omega, x).$$

It combines two results, one with Zhou Gang on equivariant graphs (surfaces of revolution), i.e.

$$u_0(\omega, x) \text{ is independent of } \omega,$$

and one in general case with Zhou Gang and Dan Knopf, ArXiv 2012.

Theorem. (Zhou Gang-S, Zhou Gang-Knopf-S) Let $d \geq 1$ and (informally for brevity)

M_0 be a surface close to a cylinder, C^{d+1} ,

M_0 has an arbitrary shallow waist and is even w.r.to the waist.

Then M_t is defined by an immersion

$$\psi(\omega, x, t) = (u(\omega, x, t)\omega, x)$$

of C^{d+1} , where $(\omega, x) \in C^{d+1}$ and $u(\omega, x, t)$ satisfies

- (i) There exists a finite time t^* such that $\inf u(\cdot, t) > 0$ for $t < t^*$ and $\lim_{t \rightarrow t^*} \inf u(\cdot, t) \rightarrow 0$;
- (ii) If $u_0 \partial_x^2 u_0 \geq -1$ then there exists a function $u_*(\omega, x) > 0$ such that $u(\omega, x, t) \geq u_*(\omega, x)$ for $\mathbb{R} \setminus \{0\}$ and $t \leq t^*$.

Theorem. (Zhou Gang-S, Zhou Gang-Knopf-S)

(iii) There exist C^1 functions $\zeta(\omega, x, t)$, $\lambda(t)$ and $b(t)$ such that

$$u(\omega, x, t) = \lambda(t) \left[\sqrt{\frac{d + b(t)y^2}{a(t)}} + \zeta(\omega, y, t) \right]$$

with $y := x/\lambda(t)$, $a(t) = -\lambda(t)\partial_t\lambda(t)$ and

$$\|\langle y \rangle^{-m} \partial_y^n \zeta(\omega, y, t)\|_\infty \leq cb^2(t), \quad m + n = 3.$$

(iv) The parameters $\lambda(t)$ and $b(t)$ satisfy (with $\tau := 2d(t^* - t)$)

$$\lambda(t) = \tau^{\frac{1}{2}}(1 + o(1)) \quad (\text{scaling eigenvalue})$$

$$b(t) = -\frac{d}{\ln \tau} \left(1 + O\left(\frac{1}{|\ln \tau|^{3/4}}\right) \right) \quad (\text{shape eigenvalue}).$$

Comparison with Previous Results

A result similar to (ii) (axi-symmetric surfaces) but for a different set of initial conditions was proven by H.M.Soner and P.E.Souganidis.

The previous result closest to ours is that by S. Angenent and D. Knopf on the axi-symmetric neckpinching for the Ricci flow.

Some ideas of the proof are close to those of Bricmont and Kupiainen on NLH.

All works mentioned above deal with *surfaces of revolution* of barbell shapes (*far from cylinders*) which are either compact (Dirichlet b.c.) or periodic (Neumann b.c.).

These works rely on parabolic maximum principle going back to Hamilton and monotonicity formulae for an entropy functional \int_{M_t} backward heat kernel $(x, t)d\mu_t$, due to Huisken and Giga and Kohn.

We explain what is going on. Rescale the MCF as

$$\varphi(u, \tau) := \lambda^{-1}(t)\psi(u, t), \quad \tau := \int_0^t \frac{dt'}{\lambda(t')^2}.$$

Important point: we do not fix $\lambda(t)$ but consider it as free parameter to be found from MCF. The rescaled MCF satisfies

$$(\partial_\tau \varphi)^N = -H(\varphi) + a\langle \varphi, \nu(\varphi) \rangle, \quad a = -\dot{\lambda}\lambda.$$

- ▶ The rescaled MCF is a gradient flow for the Huisken functional

$$V_a(\varphi) := \int_{M^\lambda} e^{-\frac{a}{2}|x|^2},$$

where $M^\lambda = \lambda^{-1}(t)M$ is the rescaled surface M .

Self-similar Surfaces

Static solutions of the rescaled MCF

$$(\partial_\tau \varphi)^N = -H(\varphi) + a\langle \varphi, \nu(\varphi) \rangle, \quad a = -\dot{\lambda} \lambda.$$

- ▶ are self-similar surfaces,

$$H(\varphi) - a\langle \nu(\varphi), \varphi \rangle = 0, \quad a \in \mathbb{R}.$$

We expect that as $\tau \rightarrow \infty$, solutions to the rescaled MCF converge to self-similar surfaces.

Hence one wants to classify self-similar surfaces and determine which ones of them are stable.

A key notion here is that of stability \Rightarrow spectrum of the hessian.

Symmetries and Spectrum of Hessian

Theorem. The hessian $\text{Hess}^N V_a(\varphi)$ of $V_a(\varphi)$ in the normal direction at a self-similar d -dimensional surface φ has

1. (Colding-Minicozzi) the simple eigenvalue $-2a$,
2. (Colding-Minicozzi) the eigenvalue $-a$ of multiplicity $d + 1$,
3. the eigenvalue 0 of multiplicity $\frac{1}{2}(d - 1)d$ (unless φ is a sphere).

These eigenvalues are due to *rescaling, translations and rotations* of the surface. The eigenvalue 0 distinguishes between a *sphere, a cylinder and a generic surface*.

Proof. Let $H_a(\varphi) := H(\varphi) - a\varphi \cdot \nu(\varphi)$. To prove say the first statement, we observe that, since $H_{\lambda^{-2}a}(\lambda\varphi) = \lambda^{-1}H_a(\varphi)$,

$$H_{\lambda^{-2}a}(\lambda\varphi) = 0, \quad \forall \lambda \in \mathbb{R}_+.$$

Differentiating this equation w.r.to λ at $\lambda = 1$, and reparametrizing the result, we arrive at the desired eigenvalue equation. \square

Spectrum and Mean convexity

The spectral information \Rightarrow the geometry of φ .

Theorem. Let φ be a self-similar surface. Then:

(a) (Colding-Minicozzi) For $a > 0$ (shrinker),

$$\text{Hess}^N V_a(\varphi) \geq -2a \text{ iff } H(\varphi) > 0.$$

(b) For $a < 0$ (expander), $H(\varphi)$ changes the sign.

Proof. The normal hessian, $\text{Hess}^N V_a(\varphi)$, has a positivity improving property

\Rightarrow the Perron-Frobenius theory applies \Rightarrow the theorem. \square

Definition. (Colding-Minicozzi) A self-similar surface is F -stable iff the normal hessian satisfies $\text{Hess}^N V_a(\varphi) \geq 0$.

Theorem. (Huisken, Colding-Minicozzi) The only self-similar, F -stable surfaces of polyn. growth are planes, spheres and cyl..

For $a = 0$, φ is a minimal surface \Rightarrow cf. Bernstein conjecture.

For a self-similar surface φ , consider the manifold of rescaled, translated and rotated self-similar surfaces

$$\mathcal{M}_{\text{self-sim}} := \{ T_{\mathbf{g}}^{\text{rot}} T_{\mathbf{z}}^{\text{transl}} T_{\lambda}^{\text{scal}} \varphi : (\lambda, \mathbf{z}, \mathbf{g}) \in \mathbb{R}_+ \times \mathbb{R}^{d+2} \times SO(d+2) \}.$$

Definition (Linearized stability of self-similar surfaces)

We say that a self-similar surface ϕ , with $a > 0$, is *linearly stable* iff

$$\text{Hess}^N V_a(\varphi) > 0 \quad \text{on} \quad \{\text{scaling, transl., rot. modes}\}^\perp.$$

(I.e. the only unstable motions allowed are scaling, transl., rot..)

The spectral theorem above gives unstable and central manifolds corresponding to the eigenvalues $-2a$, $-a$ and 0 .

Hence, if these are the only non-positive eigenvalues, then we expect the stability in the transverse direction to $\mathcal{M}_{\text{self-sim}}$. Otherwise, we expect instability.

Spectral Picture of Collapse: Sphere and Cylinder

For the d -sphere of the radius $\sqrt{\frac{a}{d}}$, the normal hessian > 0 on (scaling and translational modes) $^\perp \Rightarrow$ by the definition above, it is linearly stable.

For the $(d + 1)$ -cylinder of the radius $\sqrt{\frac{a}{d}}$, the normal hessian has, in addition to the eigenvalues above,

1. the eigenvalue $-a$ of multiplicity 1, due to translations along the axis of the cylinder,
2. the eigenvalue 0 of multiplicity $d + 1$, which originates in a "shape instability".

Hence the $(d + 1)$ -cylinder is linearly unstable.

Neckpinching

We use the eigenfunctions corresponding to instability eigenvalues,

1. due to translations along the axis of the cylinder (the eigenvalue $-a$ of multiplicity 1),
2. due to a "shape instability" (the eigenvalue 0 of multiplicity $d + 1$)

to find the approximate neck profile

$$\varphi(\omega, y, \tau) = (y, \nu_{ab}(y)\omega), \quad \text{where} \quad \nu_{ab} := \sqrt{\frac{d + by^2}{a}}, \quad b > 0.$$

Central-Unstable Manifold

Non-positive eigenvalues correspond to central-unstable manifold.
Symmetry eigenvalues correspond to the manifold of shifted and translated cylinders

$$M_{cyl} := \{\lambda(y, \omega) : (\lambda, z, g) \in G_{sym}\},$$

where $\varphi_{ab}(y, \omega) := (y, \nu_{ab}(y)\omega)$ and
 $G_{sym} := \mathbb{R} \times \mathbb{R}^{d+2} \times SO(d+2)$.

The full central-unstable manifold (the manifold of modulated cylinders or necks) is

$$M_{neck} := \{\lambda g \varphi_{ab} : (\lambda, z, g, a, b) \in \mathcal{P}\},$$

where $\varphi_{ab}(y, \omega) := (y, \nu_{ab}(y)\omega)$ and $\mathcal{P} := G_{sym} \times \mathbb{R}^+ \times \mathbb{R}^+$.

Hessian on the Neck

Consider the Hessian on the neck $\varphi_{ab} = \text{graph}_{\mathbb{C}^{d+1}} \nu_{ab}$ in the direction transversal to the neck manifold M_{neck} :

$$\text{Hess}^N V_a(\varphi_{ab}) = \underbrace{-\partial_y^2 + ay\partial_y - 2a - \frac{a}{d}\Delta_{\mathbb{S}^d}}_{\text{normal hess on cyl}} + V_{ab}(y, \omega).$$

Now, one can show that

$$\text{Hess}^N V_a(\varphi_{ab}) > 0 \quad \text{on} \quad M_{neck}^\perp$$

\Rightarrow The evolution is linearly stable in transverse directions.

Orthogonal Decomposition

We return to the original equation and look for surfaces of the form

$$\psi(x, \omega, t) = \lambda(t)g(t)\varphi(y, \omega, \tau) + z(t),$$

where $(\lambda, z, g) : [0, T) \rightarrow \mathbb{R} \times \mathbb{R}^{d+2} \times SO(d+2)$,

$$y = \lambda^{-1}(t)x, \quad \tau = \tau(t) := \int_0^t \lambda^{-2}(s)ds, \text{ and}$$

$\varphi(\cdot, \tau) := (y, u(y, \omega, \tau))$, a normal graph over the fixed cylinder.

The time dependent parameters $\lambda(t)$, $z(t)$, $g(t)$ are chosen so that $\varphi(\cdot, \tau)$ is orthogonal to the non-positive (scaling, translation and rotation) modes of the normal hessian on the cylinder.

Lemma. Given $u(y, \omega, \tau)$, there exist $a(\tau)$, $b(\tau)$ s.t.

$$u(y, \omega, \tau) = \nu_{a(\tau), b(\tau)}(y) + f(y, \omega, \tau),$$

with $f(\cdot, \tau) \perp 1$, $a(\tau)y^2 - 1$ in $L^2(\mathbb{R} \times \mathbb{S}^d, e^{-\frac{a(\tau)}{2}y^2} dyd\omega)$.

The vectors 1 and $(1 - ay^2)$ which are almost tangent vectors to the manifold, M_{neck} , provided b is sufficiently small.

Lyapunov-Schmidt Splitting (Effective Equations)

We obtained the splitting

$$u(y, \tau) = V_{a(\tau), b(\tau)}(y) + g(y, \omega, \tau) \quad (1)$$

By the the results above,

$$f(\cdot, \tau) \perp \{\text{symmetry modes}\}, \quad 1, \quad a(\tau)y^2 - 1.$$

Substitute this into (MCF) to obtain

$$\partial_\tau f = -L_{ab}f + F_{ab} + N_{ab}(f) \quad (2)$$

where $L_{ab} := \text{Hess}^N V_a(\varphi_{ab})$ is the transverse hessian, $F_{ab} \approx a$ sum of generators of broken symmetries, $N_{ab}(\phi) = \text{nonlinearity}$.

Project (2) on generators of symmetry transformations,
 $1, a(\tau)y^2 - 1 \implies$ the *equations for the parameters* a, b .

It remains to estimate f . Here we use $L_{ab} > 0$ on M_{neck}^\perp .

Let $U(\tau, \sigma)$, $\tau \geq \sigma \geq 0$, be the propagator generated by $-L_{ab}$. The main step in the proof involves showing the *key propagation estimate*: $\forall g \in X^\perp$,

$$\|\langle z \rangle^{-3} U(\tau, \sigma) g\|_\infty \lesssim e^{-c(\tau-\sigma)} \|\langle z \rangle^{-3} g\|_\infty.$$

Estimating ϕ

Let $U(\tau, \sigma)$ be the propagator generated by $-L_a$. By Duhamel principle we rewrite the differential equation for $\phi(y, \tau)$ as

$$\phi(\tau) = U(\tau, 0)\phi(0) + \int_0^\tau U(\tau, \sigma)(F + N)(\sigma)d\sigma.$$

Using the *key propagation estimate* ($\tau \geq \sigma \geq 0$)

$$\|\langle z \rangle^{-3} U(\tau, \sigma)g\|_\infty \lesssim e^{-c(\tau-\sigma)} \|\langle z \rangle^{-3}g\|_\infty,$$

where $g \perp 1$, $a(\tau)y^2 - 1$ in $L^2(\mathbb{R}, e^{-\frac{a(\tau)}{2}y^2} dy)$, we obtain

$$M(\tau) \leq M(0) + b^{\frac{1}{2}}(0)P(M(\tau)),$$

where

$$M(\tau) := \max_{\sigma \leq \tau} b^{-\frac{m+n+1}{2}}(\sigma) \|\langle y \rangle^{-m} \partial_y^n \phi(\cdot, \sigma)\|_\infty.$$

Estimating the Linear Propagator. I

Write $L_{ab} = L_{a0} + V$, with $L_{a0} := -\partial_y^2 + ay\partial_y - 2a$ (the normal hessian at the cylinder), and use that V is slowly varying in y to do a multiplicative perturbation (adiabatic) theory.

For the integral kernel $K(x, y)$ of $U(\tau, \sigma)$ (for simplicity, we do not display the variables of \mathbb{S}^d), we have the representation

$$K(x, y) = K_0(x, y)\langle e^V \rangle(x, y),$$

where $K_0(x, y)$ is the integral kernel of the operator $e^{-(\tau-\sigma)L_{a0}}$ and

$$\langle e^V \rangle(x, y) = \int e^{\int_{\sigma}^{\tau} V(\omega(s) + \omega_0(s), s) ds} d\mu(\omega).$$

Here $d\mu(\omega)$ is a harmonic oscillator (Ornstein-Uhlenbeck) probability measure on the continuous paths $\omega : [\sigma, \tau] \rightarrow \mathbb{R}$ with the boundary condition $\omega(\sigma) = \omega(\tau) = 0$ and

$$(-\partial_s^2 + a^2)\omega_0 = 0 \text{ with } \omega_0(\sigma) = y \text{ and } \omega_0(\tau) = x.$$

Estimating the Linear Propagator. II

To estimate $U(x, y)$ for $e^{a(\tau-\sigma)} \leq b^{-1/32}(\tau)$ we use the explicit formula

$$K_0(x, y) = 4\pi(1 - e^{-2ar})^{-\frac{1}{2}} \sqrt{ae^{2ar}} e^{-a \frac{(x - e^{-ary})^2}{2(1 - e^{-2ar})}},$$

where $r := \tau - \sigma$, and the bound

$$|\partial_y \langle e^V \rangle(x, y)| \leq b^{\frac{1}{2}} r,$$

which follows from the definition of $\langle e^V \rangle$ and the properties

$$V(y, \tau) \geq 0 \text{ and } |\partial_y V(y, \tau)| \lesssim b^{\frac{1}{2}}(\tau).$$

Then we iterate using the semi-group property \Rightarrow estimate of the remainder ϕ .

We do not fix the cylinder and look for surfaces of the form

$$\psi(x, \omega, t) = \lambda(t)g(t)\varphi(y, \omega, \tau) + z(t),$$

where $(\lambda, z, g) : [0, T) \rightarrow \mathbb{R} \times \mathbb{R}^{d+2} \times SO(d+2)$,
to be determined later,

$y = \lambda^{-1}(t)x$, $\tau = \tau(t) := \int_0^t \lambda^{-2}(s)ds$, and
 $\varphi(\cdot, \tau) : \mathcal{C}^{d+1} \rightarrow \mathbb{R}^{d+2}$ is a normal graph over the fixed cylinder.

The time dependent parameters $\lambda(t)$, $z(t)$, $g(t)$ are chosen so that $\varphi(\cdot, \tau)$ is orthogonal to the non-positive (scaling, translation and rotation) modes of the normal hessian on the cylinder.

Then we proceed as before.

Thank-you for your attention.

Comparison with Yang-Mills and Wave Maps Equations

Compare the dynamics for the scaling parameter $\lambda(t)$ for (MCF) and the critical Yang-Mills equation

$$\lambda \ddot{\lambda} = \frac{3}{4} \dot{\lambda}^4,$$

which gives

$$\lambda \approx \sqrt{\frac{2}{3}} \frac{t_* - t}{\sqrt{-\ln(t_* - t)}}.$$

and the critical wave map equation

$$\dot{\lambda}^2 = \lambda \ddot{\lambda} \ln \frac{a}{\lambda \ddot{\lambda}}, \quad a = 0.122.$$